

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/026157

International filing date: 12 August 2004 (12.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 10/640,904
Filing date: 14 August 2003 (14.08.2003)

Date of receipt at the International Bureau: 28 September 2006 (28.09.2006)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

September 15, 2006

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: *10/640,904*

FILING DATE: *August 14, 2003*

RELATED PCT APPLICATION NUMBER: *PCT/US04/26157*

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS *US10/640,904*



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

Transmittal of Patent Application - Rule 1.53(b)

Mail Stop Patent Application
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Inventor(s): Ross M. Kedl, George W. Griesgraber, Isidro Angelo E. Zarraga

Title: IMMUNOSTIMULATORY COMPOSITIONS AND METHODS OF STIMULATING AN IMMUNE RESPONSE

1. ☒ Enclosed is the above-identified new patent application. It includes:
 - 51 Pages of Specification (including claims and abstract)
 - 56 Number of Claims
 - 6 Sheets of Drawings
2. ☒ This application claims priority to Provisional Application No. 60/403,846, filed August 15, 2002.
3. ☒ Enclosed is an executed Declaration and Power of Attorney.
4. ☒ Enclosed is an executed Assignment to 3M Innovative Properties Company and a completed Assignment Recordation Cover Sheet.
5. ☐ Enclosed is a Certified Copy of Priority Document(s) _____ (if foreign priority is claimed).
6. ☐ Other enclosures:
7. ☒ The fee for filing this application is computed as follows:

Claims as Filed				
(1) For	(2) Number Filed	(3) Number Extra	(4) Rate	(5) Basic Fee: \$750.00
Total claims	56 - 20 =	36	x \$18.00	\$648.00
Independent claims	5 - 3 =	2	x \$84.00	\$168.00
Additional fee for filing one or more multiple dependent claims			\$280.00	\$0.00
Total Amount Due				\$1,566.00

8. ☒ Please charge to Deposit Account No. 13-3723 any fees under 37 CFR §§ 1.16 and 1.17, which may be required to file and during the entire pendency of this application. This authorization includes the fee for any necessary extension of time under 37 CFR § 1.136(a). To the extent any such extension should become necessary, it is hereby requested. One copy of this sheet marked duplicate is also enclosed.
9. ☒ Enclosed is a return receipt postcard.

Respectfully submitted,

August 14, 2003
Date

By: Christopher D. Gram
Christopher D. Gram, Reg. No.: 43,643
Telephone No.: (651) 733-1507

Office of Intellectual Property Counsel
3M Innovative Properties Company
Facsimile No.: (651) 736-3833

Filing of Papers and Fees by Express Mailing

Pursuant to 37 CFR § 1.10, this application and the documents and fees listed on this transmittal letter are being deposited on the date indicated below with the United States Postal Service "Express Mail Post Office to Addressee" service addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450

Date of Deposit <u>August 14, 2003</u>	Express Mail Label No. <u>EV326638748US</u>
---	--

- ☒ Customer Number for
Correspondence Address:



32692

PATENT TRADEMARK OFFICE

**IMMUNOSTIMULATORY COMPOSITIONS AND METHODS OF
STIMULATING AN IMMUNE RESPONSE**

Cross-Reference to Related Application

This application claims priority to U.S. Provisional Patent Application No. 60/403,846, filed August 15, 2002.

Background

Immune response modifiers ("IRMs") include compounds that possess potent immunomodulating activity including but not limited to antiviral and antitumor activity. Certain IRMs modulate the production and secretion of cytokines. For example, certain IRM compounds induce the production and secretion of cytokines such as, e.g., Type I interferons, TNF- α , IL-1, IL-6, IL-8, IL-10, IL-12, MIP-1, and/or MCP-1. As another example, certain IRM compounds can inhibit production and secretion of certain TH-2 cytokines, such as IL-4 and IL-5. Additionally, some IRM compounds are said to suppress IL-1 and TNF (U.S. Patent No. 6,518,265).

Certain IRMs are small organic molecules (e.g., molecular weight less than about 1000 Daltons, in certain cases less than about 500 Daltons, as opposed to large biologic protein, peptides, and the like) such as those disclosed in, for example, U.S. Patent Nos. 4,689,338; 4,929,624; 4,988,815; 5,037,986; 5,175,296; 5,238,944; 5,266,575; 5,268,376; 5,346,905; 5,352,784; 5,367,076; 5,389,640; 5,395,937; 5,446,153; 5,482,936; 5,693,811; 5,741,908; 5,756,747; 5,939,090; 6,039,969; 6,083,505; 6,110,929; 6,194,425; 6,245,776; 6,331,539; 6,376,669; 6,451,810; 6,525,064; 6,545,016; 6,545,017; 6,558,951; and 6,573,273; European Patent 0 394 026; U.S. Patent Publication No. 2002/0055517; and International Patent Publication Nos. WO 01/74343; WO 02/46188; WO 02/46189; WO 02/46190; WO 02/46191; WO 02/46192; WO 02/46193; WO 02/46749 WO 02/102377; WO 03/020889; WO 03/043572 and WO 03/045391.

Additional examples of small molecule IRMs include certain purine derivatives (such as those described in U.S. Patent Nos. 6,376,501, and 6,028,076), certain imidazoquinoline amide derivatives (such as those described in U.S. Patent No. 6,069,149), certain benzimidazole derivatives (such as those described in U.S. Patent 6,387,938), and certain derivatives of a 4-aminopyrimidine fused to a five membered nitrogen containing heterocyclic ring (such as adenine derivatives described in U. S. Patent Nos. 6,376,501; 6,028,076 and 6,329,381; and in WO 02/08595).

Other IRMs include large biological molecules such as oligonucleotide sequences. Some IRM oligonucleotide sequences contain cytosine-guanine dinucleotides (CpG) and are described, for example, in U.S. Patent Nos. 6,194,388; 6,207,646; 6,239,116; 6,339,068; and 6,406,705. Some CpG-containing oligonucleotides can include synthetic immunomodulatory structural motifs such as those described, for example, in U.S. Pat. Nos. 6,426,334 and 6,476,000. Other IRM nucleotide sequences lack CpG and are described, for example, in International Patent Publication No. WO 00/75304.

Certain IRMs can function as Toll-like receptor (TLR) agonists. Some small molecule IRMs may act through one or more of TLRs 2, 4, 6, 7, and 8. CpG may act through TLR 9.

By stimulating certain aspects of the immune system, as well as suppressing other aspects (see, e.g., U.S. Patent Nos. 6,039,969 and 6,200,592), IRMs may be used to treat many diseases. For example, the small molecule IRM imiquimod is useful for the treatment of external genital and perianal warts caused by human papillomavirus [see, e.g., Tomai et al, Antiviral Research 28(3): 253-64 (1995)]. Examples of other diseases that may be treated using IRMs include, but are not limited to, basal cell carcinoma, eczema, essential thrombocythaemia, hepatitis B, multiple sclerosis, neoplastic diseases, psoriasis, rheumatoid arthritis, type I herpes simplex, and type II herpes simplex.

IRM compounds also can modulate humoral immunity by stimulating antibody production by B cells. Further, various IRMs have been shown to be useful as vaccine adjuvants (see, e.g., U.S. Pat. Nos. 6,083,505 and 6,406,705).

Summary of the Invention

It has now been found that IRMs, especially small molecule IRMs and agonists of TLR 2, 4, 6, 7, and 8, are surprisingly effective at stimulating an immune response when chemically or physically paired with an antigen to form an immunostimulatory composition.

5 The immunostimulatory effect of a particular composition may be greater than the immunostimulatory effect of the same antigen and the same or a comparable IRM as that in the composition, but administered in an unpaired form.

The present invention provides an immunostimulatory composition that includes an immune response modifier (IRM) portion paired with an antigenic portion. In some
10 embodiments, the IRM portion may be, or be derived from, an agonist of Toll-like receptor 2, Toll-like receptor 4, Toll-like receptor 6, Toll-like receptor 7, or Toll-like receptor 8. In other embodiments, the IRM portion may include, or be derived from, an imidazoquinoline amine; a tetrahydroimidazoquinoline amine; an imidazopyridine amine; an aryl ether-substituted imidazopyridine amine; a 1,2-bridged imidazoquinoline amine; a 6,7-fused
15 cycloalkylimidazopyridine amine; an imidazonaphthyridine amine; a tetrahydroimidazonaphthyridine amine; an oxazoloquinoline amine; a thiazoloquinoline amine; an oxazolopyridine amine; a thiazolopyridine amine; an oxazonaphthyridine amine; or a thiazolonaphthyridine amine. In still other embodiments, the IRM portion may include, or be derived from, an organic moiety having a molecular weight of less than about 1000
20 Daltons. The antigenic portion may include an amino acid sequence, a nucleotide sequence, a lipopolysaccharide, a prion, a bacterium, a virus, or a fungus.

In another aspect, the present invention provides a method of stimulating T cells of a patient. The method includes providing an immunostimulatory composition that includes an immune response modifier paired with an antigenic portion; permitting the
25 immunostimulatory composition to bind to antigen-presenting cells, thereby activating the antigen-presenting cells; and permitting the activated antigen-presenting cells to stimulate the patient's T cells. The patient's T cells may be stimulated *in vivo* or *in vitro*.

In yet another aspect, the present invention provides a method of stimulating antibody-producing cells. The method includes providing an immunostimulatory composition that
30 comprises an immune response modifier portion paired with an antigenic portion; and

permitting the immunostimulatory composition to bind to the antibody-producing cells. The antibody-producing cells may be stimulated *in vivo* or *in vitro*.

Various other features and advantages of the present invention should become readily apparent with reference to the following detailed description, examples, claims and appended drawings. In several places throughout the specification, guidance is provided through lists of examples. In each instance, the recited list serves only as a representative group and should not be interpreted as an exclusive list.

Brief Description of the Drawings

Figure 1 shows the percentage of ovalbumin-specific activated CD8⁺ T cells generated by mice immunized with ovalbumin, as described in Example 4.

Figure 2 shows the percentage of ovalbumin-specific activated CD8⁺ T cells generated by mice immunized subcutaneously with an IRM-ovalbumin conjugate, as described in Example 4.

Figure 3 shows the percentage of ovalbumin-specific activated CD8⁺ T cells generated by mice immunized intraperitoneally with an IRM-ovalbumin conjugate, as described in Example 4.

Figure 4 shows production of interferon- γ induced by mice immunized with ovalbumin, as described in Example 5.

Figure 5 shows production of interferon- γ induced by mice immunized subcutaneously with an IRM-ovalbumin conjugate, as described in Example 5.

Figure 6 shows production of interferon- γ induced by mice immunized intraperitoneally with an IRM-ovalbumin conjugate, as described in Example 5.

Figure 7 shows an increase in the antigen-specific immune response upon providing a secondary immunization with an IRM-ovalbumin conjugate, as described in Example 6.

Figure 8 shows reduction in tumor size after immunization with a tumor-specific IRM-antigen conjugate, as described in Example 7.

Figure 9 shows expansion of activated tumor antigen-specific CD8⁺ T cells in the spleen after immunization with an IRM-tumor antigen conjugate, as described in Example 8.

Figure 10 shows expansion of activated tumor antigen-specific CD8⁺ T cells in the tumor after immunization with an IRM-tumor antigen conjugate, as described in Example 8.

Figure 11 shows the percentage of antigen presenting cells that are presenting K^b/SIINFEKL after immunization with ovalbumin, as described in Example 9.

Figure 12 the percentage of antigen presenting cells that are presenting K^b/SIINFEKL after immunization with ovalbumin plus non-conjugated IRM, as described in Example 9.

Figure 13 the percentage of antigen presenting cells that are presenting K^b/SIINFEKL after immunization with IRM-ovalbumin conjugate, as described in Example 9.

Figure 14 shows survival rates of mice immunized with ovalbumin or an IRM-ovalbumin conjugate after challenge with ovalbumin-expressing tumor cells, as described in Example 10.

Figure 15 shows expansion of antigen-specific CD8⁺ T cells after immunization with ovalbumin, as described in Example 11.

Figure 16 shows expansion of antigen-specific CD8⁺ T cells in one subject after immunization with a colloidal suspension of IRM and ovalbumin, as described in Example 11.

Figure 17 shows expansion of antigen-specific CD8⁺ T cells in a second subject after immunization with a colloidal suspension of IRM and ovalbumin, as described in Example 11.

Figure 18 shows the expansion of antigen-specific CD8⁺ T cells as a result of immunization with IRM-antigen conjugates using two different IRMs.

Figure 19 shows the fold expansion of CD8⁺ T cells over immunization with ovalbumin alone of mice immunized with various immunostimulatory compositions.

Detailed Description of Illustrative Embodiments of the Invention

The present invention provides immunostimulatory compositions (ISCs), methods of making immunostimulatory compositions, methods of eliciting an immune response using immunostimulatory compositions, and methods of enhancing the immunostimulatory activity of an IRM by pairing the IRM with another immunostimulatory component (*e.g.*, an antigen).

The ISCs may be designed to elicit a cell-mediated immune response, a humoral immune response, or both.

As noted above, many IRMs can be used as a vaccine adjuvant to increase the immune response generated against one or more antigens also presented in the vaccine. Surprisingly, certain ISCs according to the present invention can provide an even greater immune response than a vaccine containing the same or a comparable IRM and the same antigen, but in an unpaired form. In one case, an ISC provided an immune response about five-fold greater than the immune response generated by a vaccine that included the same antigen and a comparable IRM.

As used herein, the term “paired” and variations thereof refer to components associated in some chemical or physical manner so that the components are not freely dispersible from one another. For example, two components may be covalently bound to one another so that the two components are incapable of separately dispersing or diffusing. Pairing also may be achieved by, for example, non-covalent affinity binding, ionic binding, hydrophilic or hydrophobic affinity, physical entrapment, and the like. Pairing is specifically distinguished from a simple mixture of antigen and adjuvant in a conventional vaccine. In a simple mixture, the components can be free to independently disperse within the vaccinated environment. As used herein, “paired” and variations thereof confers an understanding that paired components maintain a chemical or physical association after immunization.

The immune response modifier portion may be, or be derived from, any suitable IRM. Suitable IRMs include small organic molecules, i.e., molecules having a molecular weight of less than about 1000 Daltons, although in some embodiments the IRM may have a molecular weight of less than about 700 Daltons and in some cases the IRM may have a molecular weight from about 500 Daltons to about 700 Daltons. Suitable IRMs also include agonists of one or more of TLRs 2, 4, 6, 7, 8 and 9. In some embodiments, suitable IRMs include but are not limited to the small molecule IRM compounds described above and derivatives thereof. Suitable small molecule IRMs, having a 2-aminopyridine fused to a five membered nitrogen-containing heterocyclic ring, include but are not limited to imidazoquinoline amines including but not limited to amide-substituted imidazoquinoline amines, sulfonamide-substituted imidazoquinoline amines, urea-substituted imidazoquinoline amines, aryl ether-substituted

imidazoquinoline amines, heterocyclic ether-substituted imidazoquinoline amines, amido ether-substituted imidazoquinoline amines, sulfonamido ether-substituted imidazoquinoline amines, urea-substituted imidazoquinoline ethers, and thioether-substituted imidazoquinoline amines; tetrahydroimidazoquinoline amines including but not limited to amide-substituted tetrahydroimidazoquinoline amines, sulfonamide-substituted tetrahydroimidazoquinoline amines, urea-substituted tetrahydroimidazoquinoline amines, aryl ether-substituted tetrahydroimidazoquinoline amines, heterocyclic ether-substituted tetrahydroimidazoquinoline amines, amido ether-substituted tetrahydroimidazoquinoline amines, sulfonamido ether-substituted tetrahydroimidazoquinoline amines, urea-substituted tetrahydroimidazoquinoline ethers, and thioether-substituted tetrahydroimidazoquinoline amines; imidazopyridine amines including but not limited to amide-substituted imidazopyridine amines, sulfonamido-substituted imidazopyridine amines, urea-substituted imidazopyridine amines; aryl ether-substituted imidazopyridine amines, heterocyclic ether-substituted imidazopyridine amines, amido ether-substituted imidazopyridine amines, sulfonamido ether-substituted imidazopyridine amines, urea-substituted imidazopyridine ethers, and thioether-substituted imidazopyridine amines; 1,2-bridged imidazoquinoline amines; 6,7-fused cycloalkylimidazopyridine amines; imidazonaphthyridine amines; tetrahydroimidazonaphthyridine amines; oxazoloquinoline amines; thiazoloquinoline amines; oxazolopyridine amines; thiazolopyridine amines; oxazolonaphthyridine amines; and thiazolonaphthyridine amines.

Additional suitable small molecule IRMs include the certain purine derivatives, certain imidazoquinoline amide derivatives, certain benzimidazole derivatives, and certain derivatives of a 4-aminopyrimidine fused to a five membered nitrogen containing heterocyclic ring (e.g., adenine derivatives) described above.

Other suitable IRMs include the CpGs and other IRM nucleotide sequences that lack CpG described above.

The antigenic portion can include any material that raises a cell-mediated immune response, a humoral immune response, or both. Suitable antigenic materials include but are not limited to peptides; polypeptides; lipids; glycolipids; polysaccharides; carbohydrates;

polynucleotides; prions; live or inactivated bacteria, viruses or fungi; and bacterial, viral, fungal, protozoal, tumor-derived, or organism-derived immunogens, toxins or toxoids.

Diseases for which immunostimulatory compositions of the present invention may be used as treatments include, but are not limited to:

- 5 (a) viral diseases, such as genital warts, common warts, plantar warts, hepatitis B, hepatitis C, herpes simplex virus type I and type II, molluscum contagiosum, variola, HIV, CMV, VZV, rhinovirus, adenovirus, coronavirus, influenza, para-influenza;
- (b) bacterial diseases, such as tuberculosis, and mycobacterium avium, leprosy;
- (c) other infectious diseases, such as fungal diseases, chlamydia, candida, aspergillus,
10 cryptococcal meningitis, pneumocystis carinii, cryptosporidiosis, histoplasmosis, toxoplasmosis, trypanosome infection, leishmaniasis;
- (d) neoplastic diseases, such as intraepithelial neoplasias, cervical dysplasia, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, hairy cell leukemia, Kaposi's sarcoma, melanoma, renal cell carcinoma, myelogenous leukemia, multiple myeloma, non-
15 Hodgkin's lymphoma, cutaneous T-cell lymphoma, and other cancers;
- (e) TH-2 mediated, atopic, and autoimmune diseases, such as atopic dermatitis or eczema, eosinophilia, asthma, allergy, allergic rhinitis, systemic lupus erythematosus, essential thrombocythaemia, multiple sclerosis, Ommen's syndrome, discoid lupus, alopecia areata, inhibition of keloid formation and other types of scarring, and enhancing wound healing,
20 including chronic wounds; and
- (f) as a vaccine adjuvant for use in conjunction with any material that raises either humoral and/or cell mediated immune response, such live viral and bacterial immunogens and inactivated viral, tumor-derived, protozoal, organism-derived, fungal, and bacterial immunogens, toxoids, toxins, polysaccharides, proteins, glycoproteins, peptides, cellular
25 vaccines, DNA vaccines, recombinant proteins, glycoproteins, and peptides, and the like, for use in connection with, e.g., BCG, cholera, plague, typhoid, hepatitis A, B, and C, influenza A and B, parainfluenza, polio, rabies, measles, mumps, rubella, yellow fever, tetanus, diphtheria, hemophilus influenza b, tuberculosis, meningococcal and pneumococcal vaccines, adenovirus, HIV, chicken pox, cytomegalovirus, dengue, feline leukemia, fowl plague, HSV-

1 and HSV-2, hog cholera, Japanese encephalitis, respiratory syncytial virus, rotavirus, papilloma virus, and yellow fever.

Immunostimulatory compositions of the invention include an effective amount of biological activity of both the immune response modifier portion and the antigenic portion.

5 An effective amount of biological activity of the immune response portion ("IRM activity") includes one or more of the following: an increase in cytokine production by T cells, activation of T cells specific to an antigen, and activation of dendritic cells. An effective amount of biological activity of the antigenic portion ("antigenic activity") includes one or more of the following: generation of antibodies specific to the antigen by B cells and
10 generation of antigen-presenting cells that present the antigen. Immunostimulatory compositions of the present invention may be combined with a pharmaceutically acceptable carrier, one or more excipients, or some combination of the foregoing in order to form a pharmaceutical composition.

Although the exact amount of active compound used in a pharmaceutical composition
15 of the invention will vary according to factors known to those of skill in the art, such as the physical and chemical nature of the immunostimulatory composition, the nature of the carrier, the nature of the subject's immune system (e.g., suppressed, compromised, stimulated), and the intended dosing regimen, it is anticipated that pharmaceutical compositions of the invention will contain sufficient immune response modifier portion to provide a dose of about
20 100 ng/kg to about 50 mg/kg, preferably about 10 µg/kg to about 5 mg/kg, of IRM to the subject.

A variety of dosage forms may be used, such as tablets, lozenges, capsules, parenteral formulations, syrups, creams, ointments, aerosol formulations, transdermal patches, transmucosal patches and the like.

25 Pharmaceutical compositions of the invention can be administered as the single therapeutic agent in a treatment regimen, or the pharmaceutical composition may be administered in combination with another pharmaceutical composition or with other active agents, including additional immune response modifiers, antivirals, antibiotics, antibodies, proteins, peptides, oligonucleotides, etc.

In some embodiments, the immunostimulatory immune response modifier portion may be covalently coupled to the antigenic portion to form an immunostimulatory conjugate. As used herein, “covalently coupled” refers to direct and indirect coupling of two components exclusively through covalent bonds. Direct covalent coupling may involve direct covalent binding between an atom of the immune response modifier portion and an atom of the antigenic portion. Alternatively, the covalent coupling may occur through a linking group covalently attached to the IRM portion, the antigenic portion, or both, that facilitates covalent coupling of the IRM portion and the antigenic portion. Indirect covalent coupling may include a third component such as, for example, a solid support to which the immune response modifier portion and the antigenic portion are separately covalently attached. Also, “covalently coupled” and “covalently attached” are used interchangeably.

An immunostimulatory conjugate can include an immune response modifier moiety as the IRM portion and an antigen-containing moiety as the antigenic portion. When synthesizing an immunostimulatory conjugate, each of the immune response modifier moiety, the linking group, and the antigen-containing moiety may be selected so that the resulting immunostimulatory conjugate possesses an effective amount of IRM activity and an effective amount of antigenic activity.

The linking group can be any suitable organic linking group that allows the antigen-containing moiety to be covalently coupled to the immune response modifier moiety while preserving an effective amount of IRM activity and antigenic activity. In some embodiments, the linking group may be selected to create sufficient space between the active core of the immune response modifier moiety and the antigen-containing moiety that the antigen-containing moiety does not interfere with a biologically effective interaction between the active core and T cells that results in IRM activity such as cytokine production.

The linking group includes a reactive group capable of reacting with the antigen to form a covalent bond. Suitable reactive groups include those discussed in Hermanson, G. (1996), *Bioconjugate Techniques*, Academic Press, Chapter 2 “The Chemistry of Reactive Functional Groups”, 137-166. For example, the linking group may react with a primary amine (e.g., an N-hydroxysuccinimidyl ester or an N-hydroxysulfosuccinimidyl ester); it may react with a sulfhydryl group (e.g., a maleimide or an iodoacetyl), or it may be a photoreactive

group (e.g. a phenyl azide including 4-azidophenyl, 2-hydroxy-4-azidophenyl, 2-nitro-4-azidophenyl, and 2-nitro-3-azidophenyl).

A chemically active group accessible for covalent coupling to the linking group includes groups that may be used directly for covalent coupling to the linking group or groups that may be modified to be available for covalent coupling to the linking group. For example, suitable chemically active groups include but are not limited to primary amines and sulfhydryl groups. Because certain antigen-containing moieties, e.g., proteins and other peptides, may include a plurality of chemically active groups, certain ISCs according to the present invention may include a plurality of IRM moieties conjugated to a particular antigen-containing moiety.

Methods of Making Immunostimulatory Conjugates

Immunostimulatory conjugates according to the present invention generally may be prepared by reacting an immune response modifier with a crosslinker and then reacting the resulting intermediate with an antigen. Many crosslinkers suitable for preparing bioconjugates are known and many are commercially available. See for example, Hermanson, G. (1996) *Bioconjugate Techniques*, Academic Press.


Immunostimulatory conjugates according to the present invention also may be prepared, for example, according to the method shown in Reaction Scheme I in which the antigen-containing moiety is linked to the IRM moiety through R₁. In step (1) of Reaction Scheme I a compound of Formula III is reacted with a heterobifunctional cross-linker of Formula IV to provide a compound of II. R_A and R_B each contain a functional group that is selected to react with the other. For example, if R_A contains a primary amine, then a heterobifunctional cross-linker may be selected in which R_B contains an amine-reactive functional group such as an N-hydroxysulfosuccinimidyl ester. R_A and R_B may be selected so that they react to provide the desired linker group in the conjugate.

Methods for preparing compounds of Formula III where R_A contains a functional group are known. See for example, U.S. Patent Nos. 4,689,338; 4,929,624; U.S. Patent No. 5,268,376; 5,389,640; 5,352,784; 5,494,916; 4,988,815; 5,367,076; 5,175,296; 5,395,937; 5,741,908; 5,693,811; 6,069,149; 6,194,425; and U.S. Patent 6,331,539 and International

Publications WO 00/76505; WO 00/76518; WO 02/46188, WO 02/ 46189; WO 02/46190; WO 02/46191; WO 02/46192; WO 02/46193; and WO 02/46194.

Many heterobifunctional cross-linkers are known and many are commercially available. See for example, Hermanson, G. (1996), *Bioconjugate Techniques*, Academic Press, Chapter 5 “Heterobifunctional Cross-Linkers”, 229-285. The reaction generally can be carried out by combining a solution of the compound of Formula III in a suitable solvent such as N,N-dimethylformamide with a solution of the heterobifunctional cross-linker of Formula IV in a suitable solvent such as N,N-dimethylformamide. The reaction may be run at ambient temperature. The product of Formula II may then be isolated using conventional techniques.

In step (2) of Reaction Scheme I a compound of Formula II that contains reactive group Z_A is reacted with the antigen to provide the immunostimulatory conjugate of Formula I. The reaction generally can be carried out by combining a solution of the compound of Formula II in a suitable solvent such as dimethyl sulfoxide with a solution of the antigen in a suitable buffer such as PBS. The reaction may be run at ambient temperature or at a reduced temperature (~4°C). If Z_A is a photoreactive group such as a phenyl azide then the reaction mixture will be exposed to long wave UV light for a length of time adequate to effect cross-linking (e.g., 10 – 20 minutes). The average number of immune response modifier moieties per antigen moiety may be controlled by adjusting the amount of compound of Formula II used in the reaction. The immune response conjugate of Formula I may be isolated and purified using conventional techniques.



Alternatively, a compound of Formula II may be synthesized without using a heterobifunctional cross-linker. So long as the compound of Formula II contains the reactive group Z_A , it may be reacted with the antigen using the method of step (2) above to provide an immunostimulatory conjugate.

As used herein, the terms “alkyl”, “alkenyl” and the prefix “alk-” include straight chain, branched chain, and cyclic groups, i.e. cycloalkyl and cycloalkenyl. Unless otherwise specified, these groups contain from 1 to 20 carbon atoms, with alkenyl groups containing from 2 to 20 carbon atoms. Preferred groups have a total of up to 10 carbon atoms. Cyclic groups can be monocyclic or polycyclic and preferably have from 3 to 10 ring carbon atoms. Exemplary cyclic groups include cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, and adamantyl.

The term “haloalkyl” is inclusive of groups that are substituted by one or more halogen atoms, including perfluorinated groups. This is also true of groups that include the

prefix “halo-”. Examples of suitable haloalkyl groups are chloromethyl, trifluoromethyl, and the like.

The term “aryl” as used herein includes carbocyclic aromatic rings or ring systems. Examples of aryl groups include phenyl, naphthyl, biphenyl, fluorenyl and indenyl. The term “heteroaryl” includes aromatic rings or ring systems that contain at least one ring hetero atom (e.g., O, S, N). Suitable heteroaryl groups include furyl, thienyl, pyridyl, quinolinyl, isoquinolinyl, indolyl, isoindolyl, triazolyl, pyrrolyl, tetrazolyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, benzofuranyl, benzothiophenyl, carbazolyl, benzoxazolyl, pyrimidinyl, benzimidazolyl, quinoxalinyl, benzothiazolyl, naphthyridinyl, isoxazolyl, isothiazolyl, purinyl, quinazolinyl, and so on.

“Heterocyclyl” includes non-aromatic rings or ring systems that contain at least one ring hetero atom (e.g., O, S, N) and includes all of the fully saturated and partially unsaturated derivatives of the above mentioned heteroaryl groups. Exemplary heterocyclic groups include pyrrolidinyl, tetrahydrofuranyl, morpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, thiazolidinyl, isothiazolidinyl, and imidazolidinyl.

The aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, methylenedioxy, ethylenedioxy, alkylthio, haloalkyl, haloalkoxy, haloalkylthio, halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylthio, arylalkoxy, arylalkylthio, heteroaryl, heteroaryloxy, heteroarylthio, heteroarylalkoxy, heteroarylalkylthio, amino, alkylamino, dialkylamino, heterocyclyl, heterocycloalkyl, alkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, haloalkylcarbonyl, haloalkoxycarbonyl, alkylthiocarbonyl, arylcarbonyl, heteroarylcarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, arylthiocarbonyl, heteroarylthiocarbonyl, alkanoyloxy, alkanoylthio, alkanoylamino, arylcarbonyloxy, arylcarbonythio, alkylaminosulfonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aryldiazinyl, alkylsulfonylamino, arylsulfonylamino, arylalkylsulfonylamino, alkylcarbonylamino, alkenylcarbonylamino, arylcarbonylamino, arylalkylcarbonylamino, heteroarylcarbonylamino, heteroarylalkylcarbonylamino, alkylsulfonylamino, alkenylsulfonylamino, arylsulfonylamino, arylalkylsulfonylamino, heteroarylsulfonylamino, heteroarylalkylsulfonylamino, alkylaminocarbonylamino, alkenylaminocarbonylamino,

arylaminocarbonylamino, arylalkylaminocarbonylamino, heteroarylaminocarbonylamino, heteroarylalkylaminocarbonylamino and, in the case of heterocyclyl, oxo. If other groups are described as being “substituted” or “optionally substituted”, then those groups can also be substituted by one or more of the above-enumerated substituents.

5 Certain substituents are generally preferred. For example, preferred R₂ groups include hydrogen, alkyl groups having 1 to 4 carbon atoms (i.e., methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, and cyclopropylmethyl), and alkoxyalkyl groups (e.g., methoxyethyl and ethoxymethyl). Preferably R₃ and R₄ are independently hydrogen or methyl or R₃ and R₄ join together to form a benzene ring, a pyridine ring, a 6-membered
10 saturated ring or a 6-membered saturated ring containing a nitrogen atom. One or more of these preferred substituents, if present, can be present in the compounds of the invention in any combination.

 In some embodiments, the immunostimulatory conjugates may include a solid support structure to which both the antigenic portion and the IRM portion are attached. In some
15 embodiments, the IRM portion, antigenic portion, or both may be covalently attached to the solid support using a linking group such as those described above. The solid support may include, for example, agarose beads, gold particles, and the like. The solid support may then be used to co-deliver the attached IRM portion and antigenic portion to the appropriate target cell population. Methods for attaching biomolecules to solid supports are known in the art.
20 Protocols for immobilizing biomolecules on solid supports are well known in the art and suitable reagents are available from commercial sources.

 Immunostimulatory compositions according to the present invention may contain chemical associations between the IRM portion and the antigenic portion other than covalent coupling. For example, an ISC may include an affinity interaction between the antigenic
25 portion and the IRM portion. Avidin-biotin affinity represents one example of a non-covalent interaction that may be utilized to pair an antigenic portion with an IRM portion. A biotin molecule may be chemically attached to an antigen via one of a number of functional groups present on amino acids in, for example, a proteinaceous antigen (e.g., primary amines or
30 sulfhydryl groups). An IRM portion may be conjugated to an avidin molecule by similar chemical means. The IRM portion and the antigenic portion may then be paired by the

avidin-biotin affinity interaction. Methods for biotinylating proteins and linking chemical groups to avidin are well known to one of skill in the art. Alternative affinity interactions that may be useful for making ISCs include, for example, antigen/antibody interactions, glycoprotein/lectin interactions.

5 Immunostimulatory compositions also may be formed by ionic interactions between an IRM portion and an antigenic portion. For example, an IRM portion, an antigenic portion, or both, may be chemically modified to contain oppositely charged components. The oppositely charged IRM portion and antigenic portion may then be incubated together to allow for ionic interaction between the two entities. The resulting ISC may then be
10 administered to a subject or a cell population, resulting in the co-delivery of both the IRM and the antigen to the target cells.

As in the case of covalently linked ISCs, ISCs in which the IRM portion and the antigenic portion are paired non-covalently can include a solid support.

15 **Methods of Eliciting Immune Responses Using Immunostimulatory Conjugates**

Immunostimulatory compositions according to the present invention may be used to elicit immune responses from cells of the immune system *in vitro* or *in vivo*. Thus, an ISC according to the present invention may be useful as a component of a vaccine or as an immunostimulatory factor used in *in vitro* cell culture of T cells or B cells. Indeed, IRMs may
20 be more potent immunostimulatory factors when delivered as part of an ISC according to the present invention compared to being delivered as an unpaired vaccine adjuvant. When used to elicit immune responses *in vitro*, the immune cells activated *in vitro* may be reintroduced into a patient. Alternatively, factors secreted by the activated immune cells, e.g., antibodies, cytokines, and the like, may be collected for investigative, diagnostic, and/or therapeutic uses.

25 Unless otherwise noted, a host may be immunized subcutaneously or intraperitoneally. After a sufficient time to allow the host to generate an immune response to the ISC, immune cells appropriate for the immunization site are harvested. For example, lymph nodes may be harvested from a host that had been immunized subcutaneously. Spleen cells may be harvested from a host immunized peritoneally. For some hosts, cell harvesting may include

sacrificing the hosts. In other cases, cell harvesting may include a biopsy or surgical removal of an appropriate tissue.

In one embodiment, ISCs may be used to induce the proliferation of antigen-specific T cells. A host (e.g., a mouse) may be immunized with an ISC that includes a particular antigen. After sufficient incubation in the host, certain T cells (e.g., CD8⁺ T cells) will mature to antigen-specific T cells in response to the immunization. A greater percentage of T cells will be antigen-specific in hosts immunized with ISCs compared to hosts immunized with only antigen (Figs. 1-3). The ISC may be paired by covalently coupling the IRM portion and the antigenic portion (Figs. 1-3) or, alternatively, by a non-covalent pairing such as, for example, a colloidal suspension (Figs. 15-17).

If the antigen is a protein (e.g., ovalbumin), it may not be necessary that the ISC include the entire protein. For example, an immunodominant peptide from the protein may be all that is required to induce development of T cells specific to the full protein antigen. Moreover, a booster immunization, e.g., 15 days after the initial immunization, can augment the induction of antigen-specific T cells (Fig. 7).

Immunization of a host with ISCs may be used to elicit an antigen-specific response in CD8⁺ cytotoxic T lymphocytes (CTLs). Such a response may be directed against many conditions including, but not limited to, tumors and virally-infected cell populations. ISCs also may be administered prophylactically to provide a host with a protective CTL immunity directed against future tumors or viral infections.

In another embodiment, ISCs may be used to induce cytokine production by antigen-specific T cells *in vitro*. Appropriate tissue from an immunized host may be harvested and cultured *in vitro* with antigen, thereby inducing production of one or more cytokines (e.g., IFN- γ , see Fig. 4-6). Again, in cases where the antigen is a protein, an immunodominant peptide from the antigen protein may be all that is required to induce the antigen-specific T cells in cell culture to produce and secrete the cytokines.

In another embodiment, ISCs may be used to inhibit tumor growth *in vivo*. Hosts having tumor cells expressing a particular antigen may be immunized with ISCs that contain the antigen. In some embodiments, the initial immunization may be boosted with a second immunization. Tumors harvested from hosts immunized with ISC containing the antigen

were generally smaller than the tumors harvested from hosts immunized with only the antigen (Fig. 8). Moreover, analysis of the tumors showed that tumors harvested from hosts immunized with ISC contained a higher percentage of antigen-specific T cells than tumors harvested from hosts immunized with only the antigen (Fig. 10).

5 In yet another embodiment, ISCs may be used to induce antigen-presenting cells (APCs, e.g., dendritic cells) to present peptides from the antigen within MHC class I complexes at their cell surfaces. Hosts may be immunized intravenously to induce this type of response. The response may be verified by harvesting and analyzing spleen cells for APCs presenting antigen/MHC class I complexes (Figs. 11-13).

10 In an alternative embodiment, ISCs may be used to develop antigen-specific T cells *in vitro*. For example, bone marrow cells may be harvested from a patient having a tumor that expresses a particular antigen. The harvested cells may be cultured *in vitro* with an ISC containing the antigen expressed by the tumor. Once again, if the antigen is a protein, the ISC may need only include an immunodominant peptide of the protein. Antigen-specific T cells
15 that develop *in vitro* in response to incubation with the ISC may be reintroduced to the patient.


In yet another alternative embodiment, an ISC may be administered to a subject having a tumor to increase the likelihood, duration, or both, of survival. An ISC including a tumor-specific antigenic portion administered to a mice challenged with melanoma cells
20 provided increased survival compared to mice immunized with only tumor antigen (Fig. 14).

Examples

The following examples have been selected merely to further illustrate features, advantages, and other details of the invention. It is to be expressly understood, however, that
25 while the examples serve this purpose, the particular materials and amounts used as well as other conditions and details are not to be construed in a manner that would unduly limit the scope of this invention.

Preparation of IRM Compounds

IRM Compound 1 (IRM1): *N*-[6-({2-[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-1,1-dimethylethyl}amino)-6-oxohexyl]-4-azido-2-hydroxybenzamide



5

Part A

A stirred solution of 4-chloro-3-nitroquinoline (17.3 g, 83.2 mmol) in 200 mL of anhydrous CH₂Cl₂, under N₂, was treated with triethylamine (23.2 mL, 166.4 mmol) and 1,2-diamino-2-methylpropane (9.57 mL, 91.5 mmol). After stirring overnight, the reaction mixture was diluted with 800 mL of CHCl₃ washed with H₂O (3 X 300 mL) and brine (300 mL). The organic portion was dried over Na₂SO₄ and concentrated to give 2-methyl-*N*¹-(3-nitroquinolin-4-yl)propane-1,2-diamine (21.0 g) as a bright yellow solid.

10

Part B

A solution of 2-methyl-*N*¹-(3-nitroquinolin-4-yl)propane-1,2-diamine (2.60 g, 10.0 mmol) in 50 mL of tetrahydrofuran (THF), under N₂, was cooled to 0°C and treated with 10 mL of 1N NaOH solution. Di-*tert*-butyl dicarbonate (2.18 g, 10.0 mmol) was then added to the rapidly stirred solution. The reaction mixture was then allowed to warm to ambient temperature and was stirred overnight. An additional 400 mg of di-*tert*-butyl dicarbonate was added and stirring was continued for 3 days. The reaction was then treated with ethyl acetate (200 mL) and washed with H₂O (2X) and brine. The organic portion was dried over Na₂SO₄ and concentrated to give a yellow solid that was triturated with 10% EtOAc/hexanes. The solid was isolated by filtration and dried under vacuum overnight to give *tert*-butyl 1,1-dimethyl-2-[(3-nitroquinolin-4-yl)amino]ethylcarbamate (2.80 g) as a yellow powder.

15

20

Part C

A solution of *tert*-butyl 1,1-dimethyl-2-[(3-nitroquinolin-4-yl)amino]ethylcarbamate (3.50 g, 9.72 mmol), in 150 mL of toluene was treated with 0.3 g of 5% Pt on carbon and shaken under H₂ (3 atm, 3 Kg/cm²) for 6 hours. The solution was then filtered through a Celite pad and concentrated to give 3.04 g of crude *tert*-butyl 2-[(3-aminoquinolin-4-yl)-1,1-dimethylethylcarbamate as a light orange foam.

Part D

A solution of *tert*-butyl 2-[(3-aminoquinolin-4-yl)-1,1-dimethylethylcarbamate (3.04 g, 9.21 mmol) in 50 mL of CH₂Cl₂ was cooled to 0°C and treated with triethylamine (1.41 mL, 10.13 mmol) and ethoxyacetyl chloride (1.02 mL, 10.17 mmol). After 2 hours, the reaction mixture was concentrated under reduced pressure. The resulting syrup was taken up in 100 mL of EtOH and treated with 4.5 mL of triethylamine. The solution was heated to reflux overnight. The reaction mixture was concentrated and taken up in 100 mL of CH₂Cl₂ and washed with H₂O (2X) and brine. The organic portion was dried over Na₂SO₄ and concentrated. The resulting syrup was purified by column chromatography (SiO₂, 80% EtOAc/hexanes) to give *tert*-butyl 2-[2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-1,1-dimethylethylcarbamate (1.57 g) as a peach colored foam.

Part E

A solution of *tert*-butyl 2-[2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-1,1-dimethylethylcarbamate (1.57 g, 3.94 mmol) in 30 mL of CH₂Cl₂ was treated with 3-chloroperoxybenzoic acid (77%, 1.01 g, 4.57 mmol). After stirring for 2 hours, the reaction mixture was treated with 30 mL of additional CH₂Cl₂ and was washed with 1% Na₂CO₃ solution (2 X 30 mL), H₂O and brine. The organic portion was then dried over Na₂SO₄ and concentrated to give *tert*-butyl 2-[2-(2-(ethoxymethyl)-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)-1,1-dimethylethylcarbamate (1.58 g) as a light brown foam.

Part F

A solution of *tert*-butyl 2-[2-(2-(ethoxymethyl)-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)-1,1-dimethylethylcarbamate (1.57 g, 3.79 mmol) in 20 mL of 1,2-dichloroethane was heated to 70 °C and treated with 2 mL of concentrated NH₄OH solution. To the rapidly stirred solution was added solid *p*-toluenesulfonyl chloride (795 mg, 4.17 mmol). The reaction mixture was then sealed in a pressure vessel and heating was continued for 2 hours. The reaction mixture was then cooled and treated with 50 mL of CHCl₃. The reaction mixture was then washed with H₂O, 1% Na₂CO₃ solution (3X) and brine. The organic portion was dried over Na₂SO₄ and concentrated to give the product as a light brown oil. The resulting oil was purified by column chromatography (SiO₂, 2-5% MeOH/CHCl₃) to give *tert*-butyl 2-[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-1,1-dimethylethylcarbamate (1.26 g) as a light yellow foam.

Part G

Tert-butyl 2-[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-1,1-dimethylethylcarbamate (1.26 g, 3.05 mmol) was dissolved in 10 mL of EtOH and treated with 10 mL of 2 M HCl in EtOH. After heating at reflux for 2 hours, the reaction mixture was cooled and concentrated under reduced pressure. The resulting yellow solid was dissolved in 50 mL of H₂O and extracted with CHCl₃ (20 mL). The organic layer was discarded and the aqueous portion was made basic (pH ~ 12) by addition of concentrated NH₄OH solution. This was then extracted with CHCl₃ (4 x 20 mL) and the combined organic portions were dried with Na₂SO₄ and concentrated to give 1-(2-amino-2-methylpropyl)-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinoline-4-amine (808 mg) as a light brown powder. m. p. 161.0-162.0 °C;

MS *m/z* 314 (M + H);

¹H NMR (300 MHz, d₆-DMSO) δ 8.30 (d, *J* = 7.7 Hz, 1H), 7.59 (dd, *J* = 1.2, 8.3 Hz, 1H), 7.40 (ddd, *J* = 1.0, 7.2, 8.1 Hz, 1H), 7.21 (ddd, *J* = 1.2, 7.0, 8.2 Hz, 1H), 6.57 (s, 2H), 4.94 (br s, 2H), 4.61 (br s, 2H), 3.52 (q, *J* = 7.0 Hz, 2H), 1.61 (s, 2H), 1.31 (t, *J* = 7.0 Hz, 3H), 1.07 (s, 6H);


^{13}C NMR (75 MHz, d_6 -DMSO) δ 152.4, 151.1, 145.7, 134.3, 126.8, 126.7, 121.7, 120.8, 115.7, 65.6, 65.2, 55.8, 52.5, 29.2, 15.4.
Anal. Calcd for $\text{C}_{17}\text{H}_{23}\text{N}_5\text{O}$: %C, 65.15; %H, 7.40; %N, 22.35. Found: %C, 65.04; %H, 7.52; %N, 22.07.

5

Part H

Under a nitrogen atmosphere, a solution of N-hydroxysulfosuccinimidyl (azidosalicylamido)hexanoate (100 mg, 0.204 mmol of Sulfo-LC-NHS-ASA from Pierce Biotechnology, Inc, Rockford, IL, USA) in N,N-dimethylformamide DMF, (2-4 mL) was added to a solution of 1-(2-amino-2-methylpropyl)-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinoline-4-amine (63 mg, 0.201 mmol) in DMF (5 mL). The reaction mixture was stirred under nitrogen in a foil wrapped vessel. After 2 days an additional 50 mg of N-hydroxysulfosuccinimidyl (azidosalicylamido)hexanoate was added. After about a week the reaction mixture was concentrated under reduced pressure at 55°C. The residue was partitioned between dichloromethane containing a small amount of methanol and water. The organic layer was separated and concentrated under reduced pressure. The residue was purified by flash chromatography (2 X 15 cm SiO_2 eluting with 6% methanol in chloroform) to provide 53 mg of product as a colorless glass. The glass was transferred using dichloromethane to a conical flask and then concentrated to provide a foam. The foam was dried overnight under high vacuum to provide 48 mg of a white crystalline solid. Analysis by NMR indicated the presence of dichloromethane so the material was dried overnight under high vacuum and then in a vacuum oven at 50°C for 5 hours. Analysis by HPLC indicated a purity of >93%.

IRM Compound 2 (IRM2): *N*-{6-[(2-{2-[4-amino-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethyl)amino]-6-oxohexyl}-4-azido-2-hydroxybenzamide



Part A

A solution of 2-(2-aminoethoxy)ethanol (29.0 g, 0.276 mol) in 180 mL of tetrahydrofuran (THF), under N₂, was cooled to 0°C and treated with 140 mL of 2N NaOH solution. A solution of di-*tert*-butyl dicarbonate (60.2 g, 0.276 mol) in 180 mL of THF was then added dropwise over 1 h to the rapidly stirred solution. The reaction mixture was then allowed to warm to room temperature and was stirred an additional 18 hours. The THF was then removed under reduced pressure and the remaining aqueous slurry was brought to pH 3 by addition of 150 mL of 1M H₂SO₄ solution. This was then extracted with ethyl acetate (300 mL, 100 mL) and the combined organic layers were washed with H₂O (2X) and brine. The organic portion was dried over Na₂SO₄ and concentrated to give *tert*-butyl 2-(2-hydroxyethoxy)ethylcarbamate as a colorless oil (47.1 g).

Part B

A rapidly stirred solution of *tert*-butyl 2-(2-hydroxyethoxy)ethylcarbamate (47.1 g, 0.230 mol) in 1 L of anhydrous CH₂Cl₂ was cooled to 0°C under N₂ and treated with triethylamine (48.0 mL, 0.345 mol). Methanesulfonyl chloride (19.6 mL, 0.253 mol) was then added dropwise over 30 min. The reaction mixture was then allowed to warm to room temperature and was stirred an additional 22 hours. The reaction was quenched by addition of 500 mL saturated NaHCO₃ solution and the organic layer was separated. The organic phase was then washed with H₂O (3 X 500 mL) and brine. The organic portion was dried over

Na₂SO₄ and concentrated to give 2-{2-[(*tert*-butoxycarbonyl)amino]ethoxy}ethyl methanesulfonate as a brown oil (63.5 g).

Part C

5 A stirred solution of 2-{2-[(*tert*-butoxycarbonyl)amino]ethoxy}ethyl methanesulfonate (63.5 g, 0.224 mol) in 400 mL of N,N-dimethylformamide (DMF) was treated with NaN₃ (16.1 g, 0.247 mol) and the reaction mixture was heated to 90°C under N₂. After 5 hours, the solution was cooled to room temperature and treated with 500 mL of cold H₂O. The reaction mixture was then extracted with Et₂O (3 X 300 mL). The combined
10 organic extracts were washed with H₂O (4 X 100 mL) and brine (2 X 100 mL). The organic portion was dried over MgSO₄ and concentrated to give 52.0 g of *tert*-butyl 2-(2-azidoethoxy)ethylcarbamate as a light brown oil.

Part D

15 A solution of *tert*-butyl 2-(2-azidoethoxy)ethylcarbamate (47.0 g, 0.204 mol) in MeOH was treated with 4 g of 10% Pd on carbon and shaken under H₂ (3 Kg/cm²) for 24 hours. The solution was then filtered through a Celite pad and concentrated to give 35.3 g of crude *tert*-butyl 2-(2-aminoethoxy)ethylcarbamate as a colorless liquid that was used without further purification.

20

Part E

 A stirred solution of 4-chloro-3-nitroquinoline (31.4 g, 0.151 mol) in 500 mL of anhydrous CH₂Cl₂, under N₂, was treated with triethylamine (43 mL, 0.308 mol) and *tert*-butyl 2-(2-aminoethoxy)ethylcarbamate (0.151 mol). After stirring overnight, the reaction
25 mixture was washed with H₂O (2 X 300 mL) and brine (300 mL). The organic portion was dried over Na₂SO₄ and concentrated to give a bright yellow solid. Recrystallization from ethyl acetate/hexanes gave 43.6 g of *tert*-butyl 2-{2-[(3-nitroquinolin-4-yl)amino]ethoxy}ethylcarbamate as bright yellow crystals.

Part F

A solution of *tert*-butyl 2-{2-[(3-nitroquinolin-4-yl)amino]ethoxy}ethylcarbamate (7.52 g, 20.0 mmol) in toluene was treated with 1.5 g of 5% Pt on carbon and shaken under H₂ (3 Kg/cm²) for 24 hours. The solution was then filtered through a Celite pad and concentrated to give 6.92 g of crude *tert*-butyl 2-{2-[(3-aminoquinolin-4-yl)amino]ethoxy}ethylcarbamate as a yellow syrup.

Part G

A solution of *tert*-butyl 2-{2-[(3-aminoquinolin-4-yl)amino]ethoxy}ethylcarbamate (10.2 g, 29.5 mmol) in 250 mL of anhydrous CH₂Cl₂ was cooled to 0°C and treated with triethylamine (4.18 mL, 30.0 mmol). Methoxypropionyl chloride (3.30 mL, 30.3 mmol) was then added dropwise over 5 min. The reaction was then warmed to room temperature and stirring was continued for 1 hour. The reaction mixture was then concentrated under reduced pressure to give an orange solid. This was dissolved in 250 mL of EtOH and 12.5 mL of triethylamine was added. The mixture was heated to reflux and stirred under N₂ overnight. The reaction was then concentrated to dryness under reduced pressure and treated with 300 mL of Et₂O. The mixture was then filtered and the filtrate was concentrated under reduced pressure to give a brown solid. The solid was dissolved in 200 mL of hot methanol and treated with activated charcoal. The hot solution was filtered and concentrated to give 11.1 g of *tert*-butyl 2-{2-[2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethylcarbamate as a yellow syrup.

Part H

A solution of *tert*-butyl 2-{2-[2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethylcarbamate (10.22 g, 24.7 mmol) in 250 mL of CHCl₃ was treated with 3-chloroperbenzoic acid (77%, 9.12 g, 40.8 mmol). After stirring 30 minutes, the reaction mixture was washed with 1% Na₂CO₃ solution (2 X 75 mL) and brine. The organic layer was then dried over Na₂SO₄ and concentrated to give 10.6 g of *tert*-butyl 2-{2-[2-(2-methoxyethyl)-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethylcarbamate as an orange foam that was used without further purification.

Part I

A solution of *tert*-butyl 2-{2-[2-(2-methoxyethyl)-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethylcarbamate (10.6 g, 24.6 mmol) in 100 mL of 1,2-dichloroethane was heated to 60°C and treated with 10 mL of concentrated NH₄OH solution. To the rapidly stirred solution was added solid *p*-toluenesulfonyl chloride (7.05 g, 37.0 mmol) over a 10 minute period. The reaction mixture was treated with an additional 1 mL concentrated NH₄OH solution and then sealed in a pressure vessel and heating was continued for 2 hours. The reaction mixture was then cooled and treated with 100 mL of CHCl₃. The reaction mixture was then washed with H₂O, 1% Na₂CO₃ solution (2X) and brine. The organic portion was dried over Na₂SO₄ and concentrated to give 10.6 g of *tert*-butyl 2-{2-[4-amino-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethylcarbamate as a brown foam.

Part J

Tert-butyl 2-{2-[4-amino-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethylcarbamate (10.6 g, 24.6 mmol) was treated with 75 mL of 2M HCl in ethanol and the mixture was heated to reflux with stirring. After 1.5 hours, the reaction mixture was cooled and filtered to give a gummy solid. The solid was washed ethanol and Et₂O and dried under vacuum to give the hydrochloride salt as a light brown solid. The free base was made by dissolving the hydrochloride salt in 50 mL of H₂O and treating with 10% NaOH solution. The aqueous suspension was then concentrated to dryness and the residue was treated with CHCl₃. The resulting salts were removed by filtration and the filtrate was concentrated to give 3.82 g of 1-[2-(2-aminoethoxy)ethyl]-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine as a tan powder.


MS 330 (M + H)⁺;

¹H NMR (300 MHz, DMSO-*d*₆) δ 8.10 (d, *J* = 8.1 Hz, 1 H); 7.66 (d, *J* = 8.2 Hz, 1 H); 7.40 (m, 1 H); 7.25 (m, 1 H); 6.88 (br s, 2 H); 4.78 (t, *J* = 5.4 Hz, 2 H); 3.89 (t, *J* = 4.8 Hz, 2 H); 3.84 (t, *J* = 6.9 Hz, 2 H); 3.54 (t, *J* = 5.4 Hz, 2 H); 3.31 (s, 3 H); 3.23 (t, *J* = 6.6 Hz, 2 H); 2.88 (t, *J* = 5.3 Hz, 2 H).

Part K

Under a nitrogen atmosphere, a solution of N-hydroxysulfosuccinimidyl (azidosalicylamido)hexanoate (100 mg, 0.204 mmol of Sulfo-LC-NHS-ASA from Pierce Biotechnology, Inc, Rockford, IL, USA) in DMF (2 – 4 mL) was added to a solution of 1-[2-(2-aminoethoxy)ethyl]-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (66 mg, 0.201 mmol) in DMF (5 mL). After 3.5 hours analysis by HPLC showed that no starting material was present. The reaction mixture was stirred overnight and then concentrated under reduced pressure. The residue was purified by flash chromatography (2 X 15 cm SiO₂ eluting with 8% methanol in chloroform) to provide 55 mg of product as a colorless glass. The glass was transferred using dichloromethane to a conical flask and then concentrated to provide a foam. The foam was dried overnight under high vacuum and then in a vacuum oven at 50°C for 5 hours to provide 45 mg of product as a white fluffy solid. Analysis by HPLC indicated a purity of >95%.

IRM Compound 3 (IRM3): *N*-(2-{2-[4-amino-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethyl)hexadecanamide



Under a nitrogen atmosphere, a suspension of 1-[2-(2-aminoethoxy)ethyl]-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (140.5 mg, 0.428 mmol) in a mixture of

dichloromethane (3.5 mL) and triethylamine (150 μ L, 1.07 mmol) was cooled to 0 °C. Palmitoyl chloride (130 μ L, 0.428 mmol) was slowly added. The reaction mixture was allowed to stir at 0 °C for 2 hours at which time analysis by thin layer chromatography indicated that there was no starting material left. The reaction mixture was diluted with
5 dichloromethane (30 mL), washed with saturated sodium bicarbonate solution (2 x 5 mL), dried over magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (12 g of silica gel eluted with 2% methanol in dichloromethane) to provide 183 mg of *N*-(2-{2-[4-amino-2-(2-methoxyethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]ethoxy}ethyl)hexadecanamide as a white powder.

10 Anal. Calcd for $C_{33}H_{53}N_5O_3$: %C, 69.80; %H, 9.41; %N, 12.33; Found: %C, 69.60; %H, 9.28; %N, 11.99.

Example 1: Crosslinking of Immune Response Modifier to Ovalbumin

IRM1 was suspended in dimethyl sulfoxide (DMSO) to 10 mg/ml. Ovalbumin was
15 suspended in phosphate buffered saline (PBS) to 10 mg/ml and the pH adjusted to > 10.0 by the addition of NaOH. 500 μ L of the ovalbumin solution (5 mg ovalbumin) was mixed with 100 μ L of the IRM1 solution (1 mg IRM1) in a single well of a 12-well tissue culture plate. The plate was placed on ice and a long wavelength UV light source was placed directly over the plate as close to the well containing the IRM1/ovalbumin mixture as possible. The
20 mixture was irradiated for 15 minutes. The resulting conjugate was removed from the well and resuspended in PBS to a final concentration of 5 mg/ml ovalbumin, 0.5 mg/ml IRM1, and dialyzed against PBS to remove any unconjugated IRM.

Example 2: Immunizations

25 C57BL/6 mice were immunized with conjugate (1 mg ovalbumin and 200 μ g IRM1, prepared as in Example 1) in 200 μ L PBS either subcutaneously or intraperitoneally. Control mice were immunized with 1 mg ovalbumin in 200 μ L PBS. For analysis of primary responses, mice were sacrificed 5-7 days after immunization. For analysis of secondary responses, the mice were boosted 7-15 days after the initial immunization and sacrificed 5-7
30 days later. Unless otherwise indicated, lymph nodes were harvested from mice immunized

subcutaneously for analysis and spleen cells were harvested from mice immunized intraperitoneally for analysis.

Example 3: Reagents

5 Fluorochrome-labeled antibodies specific for mouse CD8, CD11c, and CD44 were obtained from Pharmingen (San Diego, CA). The monoclonal antibody 25D1.16 (Dr. Ron Germain, NIH) is specific for the dominant ovalbumin peptide (SIINFEKL) bound to the MHC class I molecule H-2K^b. Porgador et al., *Immunity* 6:715-26. Ovalbumin was obtained from Sigma Chemical Company (St. Louis, MO). Tetramers of the MHC class I molecule H-2K^b bound to the dominant ovalbumin peptide SIINFEKL were produced as described in Kedl et al., *J Exp Med*, 192:1105-13 (2000). The ovalbumin-expressing melanoma cell line B16ova was made by lipofection of the B16-F10 cell line (ATCC # CRL-6475) with a plasmid encoding full-length ovalbumin. See Kedl et al., *Proc. Natl. Acad. Sci. USA*, 98:10811-6. Cytokine capture and detection reagents were from Miltenyi Biotech (Auburn, CA).

Example 4

Experimental mice were immunized with conjugate as prepared in Example 1 either subcutaneously or intraperitoneally as described in Example 2. Control mice were immunized with ovalbumin as described in Example 2. Seven days later, lymph nodes or spleen were removed and the cells were stained with antibodies specific for CD8, CD44, and H-2K^b/SIINFEKL tetramers and analyzed by flow cytometry. The results are shown in Figures 1-3. The cells that stain positively for all three markers identify activated, ovalbumin-specific CD8⁺ T cells that developed as a result of the immunization.

Each of Figures 1-3 includes the percentage of all CD8⁺ T cells examined that were identified as activated, ovalbumin-specific CD8⁺ T cells. In the control mice, 0.07% of CD8⁺ T cells were activated, ovalbumin-specific CD8⁺ T cells. Subcutaneous immunization with conjugate generated 0.52% activated, ovalbumin-specific CD8⁺ T cells. Intraperitoneal immunization with conjugate generated 0.92% activated, ovalbumin-specific CD8⁺ T cells.

Immunization with the dominant ovalbumin peptide (SIINFEKL) conjugated to IRM1 (100 µg peptide crosslinked to 200 µg IRM1) also induced the activation of ovalbumin-specific CD8⁺ T cells.

5 **Example 5**

Control mice were immunized with ovalbumin as described in Example 2. Experimental mice were immunized with conjugate as prepared in Example 1 either subcutaneously or intraperitoneally as described in Example 2. Appropriate immune cells were harvested seven days after immunization and incubated *in vitro* for four hours with
10 SIINFEKL peptide. Interferon gamma (IFN-γ)-producing CD8⁺ T cells were identified using an IFN-γ capture and detection assay coupled with flow cytometry. CD8⁺ T cells that were generated by immunizing with the IRM1-ovalbumin conjugate produced IFN-γ in response to subsequent antigen stimulation (Figures 4-6).

15 **Example 6**

Mice were immunized with conjugate as prepared in Example 1 as described in Example 2 and then boosted with an equal amount of conjugate on day 15. The spleen and lymph node cells from the boosted animals were analyzed seven days after boosting. Analysis showed an increased percentage of activated ovalbumin-specific CD8⁺ T cells over
20 those from mice that had received only the primary immunization (Figure 7).

Example 7

Mice were injected intradermally with the ovalbumin-expressing melanoma cell line B16ova (1 x 10⁵ cells). On day 7, the mice were injected subcutaneously with either 1 mg
25 ovalbumin (control) or with conjugate as prepared in Example 1. On Day 14, the mice were boosted subcutaneously with ovalbumin or conjugate as on Day 7. On Day 20, tumor size was measured with calipers in two dimensions. Immunization with the conjugate resulted in reduced tumor size as compared to the control (Figure 8).

Example 8

Mice were injected intradermally with the ovalbumin-expressing melanoma cell line B16ova (1×10^5 cells). Seven and 14 days later, the mice were injected intraperitoneally with either 1 mg ovalbumin (control) or with conjugate as prepared in Example 1. Twenty days after challenge with the melanoma cell line, spleens and tumors were removed and the cells from each source were stained with antibodies specific for CD8 and CD44 as well as H-2K^b/SIINFEKL tetramers. Flow cytometric analysis revealed a significant expansion of activated ovalbumin-specific CD8⁺ T cells both in the spleen (Figure 9) and within the tumor (Figure 10).

Example 9

Mice were immunized intravenously with ovalbumin, ovalbumin and unpaired IRM1: 1-(2-amino-2-methylpropyl)-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinoline-4-amine (IRM1 prior to attachment of the linker group – the product of Part G of the synthesis of IRM1), or with conjugate as prepared in Example 1. Fourteen hours later, the spleens were removed, collagenase treated, and the cells were stained with antibodies specific for CD8, CD11c, and the antibody 25D1.16, which is specific for the ovalbumin peptide SIINFEKL complexed with the MHC class I H-2K^b molecule. Flow cytometric analysis indicated that immunizing mice with the IRM1-ovalbumin conjugate (Figure 13) generated a greater percentage of CD11c⁺, CD8⁺ dendritic cells that were presenting K^b/SIINFEKL than immunizing mice with ovalbumin alone (Figure 11) or a mixture of unpaired ovalbumin and IRM1 (Figure 12).

Example 10

Mice were immunized on Day 0 subcutaneously as described in Example 2 with ovalbumin (control) or with IRM1 conjugate as prepared in Example 1. The mice received boost immunizations on Day 14. The mice were then challenged intradermally with the ovalbumin-expressing melanoma cell line B16ova (1×10^5 cells) on Day 28 and monitored for tumor growth. At Day 75, 80% of the mice immunized with conjugate survived and appeared healthy, while all of the mice immunized with only ovalbumin died (Fig 14).

Example 11

A stock solution of IRM3 was prepared by dissolving IRM3 in DMSO to a concentration of 10 mg/ml. Ovalbumin was dissolved in PBS to a concentration of 50 mg/ml. 50 µl of the IRM3 stock solution was added to 150 µl of PBS and then mixed by vortexing. 50 µl of the ovalbumin was added to the IRM3 solution and mixed by vortexing. A cloudy colloidal suspension of IRM3 and ovalbumin resulted.

Mice were immunized on Day 0 subcutaneously as described in Example 2 with either (a) ovalbumin alone, or (b) 50 µl of the colloidal suspension of ovalbumin and IRM3. On Day 6, draining lymph nodes were removed, homogenized, and stained with the H-2K^b/SIINFEKL tetramer to identify ovalbumin-specific T cells. Figure 15 shows flow cytometry data from a control mouse immunized with ovalbumin alone; Figures 16 and 17 show data from two different mice that were immunized with the colloidal suspension.

Example 12

A conjugate of IRM2 and ovalbumin was prepared as described in Example 1, except that IRM2 was used in place of IRM1.

Twenty-four mice were divided into eight groups of three. Four groups of three mice were immunized subcutaneously as described in Example 2 at Day 0, with each group receiving increasing amounts of IRM1-ovalbumin conjugate. The remaining four groups were similarly immunized at Day 0 with the IRM2-ovalbumin conjugate. At day 6, the mice were sacrificed and draining lymph nodes were removed and stained with the H-2K^b/SIINFEKL tetramer to identify ovalbumin-specific T cells. The percentage of ovalbumin-specific T cells was calculated for each mouse. Figure 18 summarizes the results.

Example 13

Mice were adoptively transferred with approximately 2×10^6 OT1 ovalbumin-specific transgenic T cells (The Jackson Laboratory, Bar Harbor, Maine). Mice were immunized on Day 0 subcutaneously as described in Example 2 with either (a) 100 µg ovalbumin, (b) 100 µg ovalbumin + 50 µg CpG (Ova + CpG), (c) 100 µg ovalbumin + 50 µg unpaired IRM1 (Ova + IRM), or (d) 100 µg ovalbumin conjugated to 50 µg IRM1 (Ova x IRM). On Day 5, draining

lymph nodes were removed and the cells were stained with the H-2K^b/SIINFEKL tetramer to identify ovalbumin-specific T cells. Figure 19 shows the fold expansion of ovalbumin-specific T cells in the Ova + CpG, Ova + IRM, and Ova x IRM groups over the mice immunized with ovalbumin alone.

5 The complete disclosures of the patents, patent documents and publications cited herein are incorporated by reference in their entirety as if each were individually incorporated. In case of conflict, the present specification, including definitions, shall control.

10 Various modifications and alterations to this invention will become apparent to those skilled in the art without departing from the scope and spirit of this invention. Illustrative embodiments and examples are provided as examples only and are not intended to limit the scope of the present invention. The scope of the invention is limited only by the claims set forth as follows.

What is Claimed is:

1. An immunostimulatory composition comprising:

an immune response modifier portion paired with an antigenic portion..

5 2. The immunostimulatory composition of claim 1 wherein the immune response modifier portion is an agonist of Toll-like receptor 2, Toll-like receptor 4, Toll-like receptor 6, Toll-like receptor 7, or Toll-like receptor 8.

10 3. The immunostimulatory composition of claim 1 wherein the immune response modifier portion comprises an imidazoquinoline amine; a tetrahydroimidazoquinoline amine; an imidazopyridine amine; an aryl ether-substituted imidazopyridine amine; a 1,2-bridged imidazoquinoline amine; a 6,7-fused cycloalkylimidazopyridine amine; an imidazonaphthyridine amine; a tetrahydroimidazonaphthyridine amine; an oxazoloquinoline amine; a thiazoloquinoline amine; an oxazolopyridine amine; a thiazolopyridine amine; an
15 oxazolonaphthyridine amine; or a thiazolonaphthyridine amine.

4. The immunostimulatory composition of claim 1 wherein the immune response modifier portion comprises an organic moiety having a molecular weight of less than about
20 1000 Daltons.

5. The immunostimulatory composition of claim 1 wherein immune response modifier portion and the antigenic portion are covalently coupled.

25 6. The immunostimulatory composition of claim 1 wherein the immune response modifier portion and the antigenic portion are paired by a physical or chemical association other than covalent coupling that limits independent diffusion of the immune response modifier portion with respect to the antigenic portion.


7. The immunostimulatory composition of claim 1 wherein the composition comprises a colloidal suspension.

8. The immunostimulatory composition of claim 1 wherein the antigenic portion comprises an amino acid sequence, a nucleotide sequence, a lipopolysaccharide, a prion, a bacterium, a virus, or a fungus.

9. The immunostimulatory composition of claim 8 wherein the amino acid sequence is a polypeptide.

10. The immunostimulatory composition of claim 9 wherein the polypeptide is a protein.

11. The immunostimulatory composition of claim 1 wherein the immune response modifier portion is a compound of the formula:



wherein:

R₁ is a linker group;

R₂ is selected from the group consisting of:

-hydrogen;

-alkyl;

-alkenyl;

-aryl;

-substituted aryl;

-heteroaryl;

	-substituted heteroaryl;
	-alkyl-O-alkyl;
	-alkyl-S-alkyl;
	-alkyl-O-aryl;
5	-alkyl-S-aryl;
	-alkyl-O- alkenyl;
	-alkyl-S- alkenyl; and
	-alkyl or alkenyl substituted by one or more substituents selected from the group consisting of:
10	-OH;
	-halogen;
	-N(R ₅) ₂ ;
	-CO-N(R ₅) ₂ ;
	-CS-N(R ₅) ₂ ;
15	-SO ₂ -N(R ₅) ₂ ;
	-NR ₅ -CO-C ₁₋₁₀ alkyl;
	-NR ₅ -CS-C ₁₋₁₀ alkyl;
	-NR ₅ - SO ₂ -C ₁₋₁₀ alkyl;
	-CO-C ₁₋₁₀ alkyl;
20	-CO-O-C ₁₋₁₀ alkyl;
	-N ₃ ;
	-aryl;
	-substituted aryl;
	-heteroaryl;
25	-substituted heteroaryl;
	-heterocyclyl;
	-substituted heterocyclyl;
	-CO-aryl;
	-CO-(substituted aryl);
30	-CO-heteroaryl; and

-CO-(substituted heteroaryl);

R₃ and **R₄** are each independently:

-hydrogen;

-halogen;

5 -alkyl;

-alkenyl;

-O-alkyl;

-S-alkyl; and

-N(R₅)₂;

10 or when taken together, **R₃** and **R₄** form a fused aryl or heteroaryl group that is optionally substituted by one or more substituents selected from the group consisting of;

-halogen;

-alkyl;

15 -alkenyl;

-O-alkyl;

-S-alkyl; and

-N(R₅)₂;

20 or when taken together, **R₃** and **R₄** form a fused 5 to 7 membered saturated ring, optionally containing one or more heteroatoms and optionally substituted by one or more substituents selected from the group consisting of;

-halogen;

-alkyl;

-alkenyl;


25 -O-alkyl;

-S-alkyl; and

-N(R₅)₂; and

each **R₅** is independently hydrogen or C₁₋₁₀ alkyl.

30 12. An immunostimulatory conjugate of the formula:



wherein:

- 5 **R₁** is a linker group;
- R₂** is selected from the group consisting of:
- hydrogen;
 - alkyl;
 - alkenyl;
 - 10 -aryl;
 - substituted aryl;
 - heteroaryl;
 - substituted heteroaryl;
 - alkyl-O-alkyl;
 - 15 -alkyl-S-alkyl;
 - alkyl-O-aryl;
 - alkyl-S-aryl;
 - alkyl-O- alkenyl;
 - alkyl-S- alkenyl; and
 - 20 -alkyl or alkenyl substituted by one or more substituents selected from the
 - group consisting of:
 - OH;
 - halogen;
 - N(R₅)₂;
 - 25 -CO-N(R₅)₂;
 - CS-N(R₅)₂;

5 -SO₂-N(R₅)₂;
 -NR₅-CO-C₁₋₁₀ alkyl;
 -NR₅-CS-C₁₋₁₀ alkyl;
 -NR₅-SO₂-C₁₋₁₀ alkyl;
 -CO-C₁₋₁₀ alkyl;
 -CO-O-C₁₋₁₀ alkyl;
 -N₃;
 -aryl;
 -substituted aryl;
 10 -heteroaryl;
 -substituted heteroaryl;
 -heterocyclyl;
 -substituted heterocyclyl;
 -CO-aryl;
 15 -CO-(substituted aryl);
 -CO-heteroaryl; and
 -CO-(substituted heteroaryl);

R₃ and R₄ are each independently:

20 -hydrogen;
 -halogen;
 -alkyl;
 -alkenyl;
 -O-alkyl;
 -S-alkyl; and
 25 -N(R₅)₂;

or when taken together, R₃ and R₄ form a fused aryl or heteroaryl group that is optionally substituted by one or more substituents selected from the group consisting of;

30 -halogen;
 -alkyl;

-alkenyl;
-O-alkyl;
-S-alkyl; and
-N(R₅)₂;

5 or when taken together, R₃ and R₄ form a fused 5 to 7 membered saturated ring, optionally containing one or more heteroatoms and optionally substituted by one or more substituents selected from the group consisting of;

-halogen;
-alkyl;
10 -alkenyl;
-O-alkyl;
-S-alkyl; and
-N(R₅)₂;

each R₅ is independently hydrogen or C₁₋₁₀ alkyl; and

15 n is 1 to 10; or

a pharmaceutically acceptable salt thereof.

13. The immunostimulatory conjugate of claim 12 wherein R₂ is selected from the group consisting of hydrogen, alkyl and alkyl-O-alkyl.

20

14. The immunostimulatory conjugate of claim 13 wherein R₂ is selected from the group consisting of hydrogen, methyl, ethyl, propyl, butyl, cyclopropylmethyl, ethoxymethyl and methoxyethyl.

25 15. The immunostimulatory conjugate of claim 12 wherein R₃ and R₄ are taken together to form a benzene ring.

16. The immunostimulatory conjugate of claim 12 wherein R₃ and R₄ are taken together to form a pyridine ring.


30

17. The immunostimulatory conjugate of claim 12 wherein R₃ and R₄ are taken together to form a 6-membered saturated ring.

18. The immunostimulatory conjugate of claim 12 wherein R₃ and R₄ are taken together to form a 6-membered saturated ring containing a nitrogen atom.

19. The immunostimulatory conjugate of claim 12 wherein R₃ and R₄ are independently selected from the group consisting of hydrogen and alkyl.

20. The immunostimulatory conjugate of claim 12 wherein R₁ has the formula:



wherein

X is an alkylene group optionally interrupted by one or more of:

-O-;

-S(O)₀₋₂-;

-N(R₅)-;

-N(R₅)-C(O)-;

-N(R₅)-S(O₂)-;

-N(R₅)-C(O)-N(R₅)-;

-N(R₅)-C(S)-N(R₅)-; and

-C(O)-N(R₅)-;

Y is a bond or is selected from the group consisting of:

-N(R₅)-;

-N(R₅)-C(O)-;

-N(R₅)-C(O)-cyclohexyl-;

-N(R₅)-C(O)-phenylene-; and

-phenylene-;

Z is the divalent linking group that is formed when the reactive group **Z_A** covalently couples to the antigen;

Z_A is a reactive group that reacts with the antigen to form a covalent bond; and

each R_5 is independently hydrogen or C_{1-10} alkyl.

21. The immunostimulatory conjugate of claim 20 wherein Z_A is a group that reacts with primary amines.

5

22. The immunostimulatory conjugate of claim 21 wherein Z_A is an N-hydroxysuccinimide ester or an N-hydroxysulfosuccinimide ester.

10

23. The immunostimulatory conjugate of claim 20 wherein Z_A is a group that reacts with a sulfhydryl group.

24. The immunostimulatory conjugate of claim 23 wherein Z_A is a maleimide or an iodoacetyl group.

15

25. The immunostimulatory conjugate of claim 20 wherein Z_A is a photoreactive group.

26. The immunostimulatory conjugate of claim 25 wherein Z_A is a phenyl azide.

20


27. The immunostimulatory conjugate of claim 26 wherein Z_A is selected from the group consisting of 4-azidophenyl, 2-hydroxy-4-azidophenyl, 2-nitro-4-azidophenyl, and 2-nitro-3-azidophenyl.

28. The immunostimulatory conjugate of claim 20 wherein Z is selected from the group consisting of:

25


-C(O)-;

-C(O)-CH₂-;



wherein R' is hydrogen, hydroxy or nitro.

- 5 29. The immunostimulatory conjugate of claim 12 wherein the antigen comprises an amino acid sequence, a nucleotide sequence or a lipopolysaccharide.
30. The immunostimulatory conjugate of claim 29 wherein the amino acid sequence is a polypeptide.
- 10 31. The immunostimulatory conjugate of claim 30 wherein the polypeptide is a protein.
32. A compound of the formula:



wherein:

X is an alkylene group optionally interrupted by one or more of:

- O-;
- 20 -S(O)₀₋₂-;
- N(R₅)-;

-N(R₅)-C(O)-;
-N(R₅)-S(O₂)-;
-N(R₅)-C(O)-N(R₅)-;
-N(R₅)-C(S)-N(R₅)-; and
-C(O)-N(R₅)-;

Y is a bond or is selected from the group consisting of:

-N(R₅)-;
-N(R₅)-C(O)-;
-N(R₅)-C(O)-cyclohexyl-;
-N(R₅)-C(O)-phenylene-; and
-phenylene-;

Z_A is a reactive group that reacts with the antigen to form a covalent bond;

R₂ is selected from the group consisting of:

-hydrogen;
-alkyl;
-alkenyl;
-aryl;
-substituted aryl;
-heteroaryl;
-substituted heteroaryl;
-alkyl-O-alkyl;
-alkyl-S-alkyl;
-alkyl-O-aryl;
-alkyl-S-aryl;
-alkyl-O- alkenyl;
-alkyl-S- alkenyl; and
-alkyl or alkenyl substituted by one or more substituents selected from the
group consisting of:

-OH;
-halogen;

5 -N(R₅)₂;
 -CO-N(R₅)₂;
 -CS-N(R₅)₂;
 -SO₂-N(R₅)₂;
 -NR₅-CO-C₁₋₁₀ alkyl;
 -NR₅-CS-C₁₋₁₀ alkyl;
 -NR₅-SO₂-C₁₋₁₀ alkyl;
 -CO-C₁₋₁₀ alkyl;
 -CO-O-C₁₋₁₀ alkyl;
 10 -N₃;
 -aryl;
 -substituted aryl;
 -heteroaryl;
 -substituted heteroaryl;
 15 -heterocyclyl;
 -substituted heterocyclyl;
 -CO-aryl;
 -CO-(substituted aryl);
 -CO-heteroaryl; and
 20 -CO-(substituted heteroaryl);

R₃ and **R₄** are each independently:

25 -hydrogen;
 -halogen;
 -alkyl;
 -alkenyl;
 -O-alkyl;
 -S-alkyl; and
 -N(R₅)₂;

or when taken together, R₃ and R₄ form a fused aryl or heteroaryl group that is optionally substituted by one or more substituents selected from the group consisting of;

-halogen;
-alkyl;
-alkenyl;
-O-alkyl;
-S-alkyl; and
-N(R₅)₂;

or when taken together, R₃ and R₄ form a fused 5 to 7 membered saturated ring, optionally containing one or more heteroatoms and optionally substituted by one or more substituents selected from the group consisting of;

-halogen;
-alkyl;
-alkenyl;
-O-alkyl;
-S-alkyl; and
-N(R₅)₂;

each R₅ is independently hydrogen or C₁₋₁₀ alkyl; or
a pharmaceutically acceptable salt thereof.

33. A method of stimulating T cells of a patient, the method comprising:

- a) providing an immunostimulatory composition that comprises an immune response modifier portion paired with an antigenic portion;
- b) permitting the immunostimulatory composition to bind to antigen-presenting cells, thereby activating the antigen-presenting cells; and
- c) permitting the activated antigen-presenting cells to stimulate the patient's T cells.

34. The method of claim 33 wherein the immune response modifier portion is an agonist of Toll-like receptor 2, Toll-like receptor 4, Toll-like receptor 6, Toll-like receptor 7, or Toll-like receptor 8.

35. The method of claim 33 wherein the immune response modifier portion comprises an imidazoquinoline amine; a tetrahydroimidazoquinoline amine; an imidazopyridine amine; an aryl ether-substituted imidazopyridine amine; a 1,2-bridged imidazoquinoline amine; a 6,7-fused cycloalkylimidazopyridine amine; an imidazonaphthyridine amine; a tetrahydroimidazonaphthyridine amine; an oxazoloquinoline amine; a thiazoloquinoline amine; an oxazolopyridine amine; a thiazolopyridine amine; an oxazonaphthyridine amine; or a thiazolonaphthyridine amine.

36. The method of claim 33 wherein the immune response modifier portion and the antigenic portion are covalently coupled.

37. The method of claim 33 wherein the immune response modifier portion and the antigenic portion are paired by a chemical or physical association other than covalent coupling that limits independent diffusion of the immune response modifier portion with respect to the antigenic portion.

38. The method of claim 37 wherein the immunostimulatory composition comprises a colloidal suspension.

39. The method of claim 33 wherein the immunostimulatory composition is permitted to bind to the antigen-presenting cells *in vitro*.

40. The method of claim 39 wherein the step of permitting the immunostimulatory composition to bind to antigen-presenting cells comprises:

- a) providing a cell culture comprising antigen-presenting cells; and

b) contacting an amount of the immunostimulatory composition effective for stimulating the antigen-presenting cells with the cell culture.

41. The method of claim 39 wherein the step of permitting the activated antigen-presenting cells to stimulate the patient's T cells comprises injecting the activated antigen-presenting cells into the patient.

42. The method of claim 33 wherein the immunostimulatory composition is permitted to bind to the antigen-presenting cells *in vivo*.

43. The method of claim 42 wherein the step of permitting the immunostimulatory composition to bind to antigen-presenting cells comprises:

a) providing the immunostimulatory composition as a component of a vaccine; and

b) vaccinating the patient with the vaccine.

44. The method of claim 42 wherein the immunostimulatory composition comprises a tumor-specific antigenic portion.

45. A method of stimulating antibody-producing cells, the method comprising:

a) providing an immunostimulatory composition that comprises an immune response modifier portion paired with an antigenic portion; and

b) permitting the immunostimulatory composition to bind to the antibody-producing cells.

46. The method of claim 45 wherein the immune response modifier portion comprises an agonist of Toll-like receptor 2, Toll-like receptor 4, Toll-like receptor 6, Toll-like receptor 7, or Toll-like receptor 8.

47. The method of claim 45 wherein the immune response modifier portion comprises an imidazoquinoline amine; a tetrahydroimidazoquinoline amine; an imidazopyridine amine; an aryl ether-substituted imidazopyridine amine; a 1,2-bridged imidazoquinoline amine; a 6,7-fused cycloalkylimidazopyridine amine; an imidazonaphthyridine amine; a
5 tetrahydroimidazonaphthyridine amine; an oxazoloquinoline amine; a thiazoloquinoline amine; an oxazolopyridine amine; a thiazolopyridine amine; an oxazolonaphthyridine amine; or a thiazolonaphthyridine amine.

48. The method of claim 45 wherein the immune response modifier portion and the
10 antigenic portion are covalently coupled.

49. The method of claim 45 wherein the immune response modifier portion and the antigenic portion are paired by a chemical or physical association other than covalent coupling that limits independent diffusion of the immune response modifier portion with
15 respect to the antigenic portion.

50. The method of claim 49 wherein the immunostimulatory composition comprises a colloidal suspension.

51. The method of claim 45 wherein the immunostimulatory composition is permitted to bind to the antibody-producing cells *in vivo*.

52. The method of claim 51 wherein the step of permitting the immunostimulatory composition to bind to antibody-producing cells comprises:

- 25 a) providing the immunostimulatory composition as a component of a vaccine; and
b) introducing the vaccine into a patient having antibody-producing cells.

53. The method of claim 51 wherein the immunostimulatory composition comprises a
30 tumor-specific antigenic portion.

54. The method of claim 45 wherein the immunostimulatory composition is permitted to bind to the antibody-producing cells *in vitro*.

5 55. The method of claim 54 wherein the step of permitting the immunostimulatory composition to bind to the antibody-stimulating cells comprises:

- a) providing a cell culture comprising antibody-producing cells;
- b) contacting the cell culture with an amount of the immunostimulatory composition effective for stimulating the antibody-producing cells.

10

56. The method of claim 45 further comprising collecting antibodies produced by the stimulated antibody-producing cells.

**IMMUNOSTIMULATORY COMPOSITIONS AND METHODS OF STIMULATING
AN IMMUNE RESPONSE**

Abstract of the Disclosure

- 5 The present invention provides immunostimulatory compositions that include an immune response modifier portion paired with an antigenic portion.

1/6




FIG. 1




FIG. 2




FIG. 3

2/6




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14

5/6




FIG. 15




FIG. 16




FIG. 17

6/6





FIG. 19

DECLARATION AND POWER OF ATTORNEY, FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR § 1.63)	Case No:	57971US011
	First Named Inventor:	Ross M. Kedl
	COMPLETE IF KNOWN	
	Application No.:	
	Filing Date:	
	Art Unit:	
<input checked="" type="checkbox"/> Declaration Submitted with Initial Filing <input type="checkbox"/> Declaration Submitted after Initial Filing (surcharge 37 CFR § 1.16(e) required)		Examiner Name:

As a below named inventor, I hereby declare that my residence, mailing address, and citizenship are as stated below next to my name, and that I believe I am an original, first, and sole inventor (if only one name is listed below) or an original, first, and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**IMMUNOSTIMULATORY COMPOSITIONS AND METHODS OF
STIMULATING AN IMMUNE RESPONSE**

The specification of which

☒ is attached hereto;

☐ was filed on _____
As United States Application No. _____

☐ is identified as PCT International Application No. _____
filed on _____

and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 USC §§ 119(a)-(d) or (f), or 365(b) of any foreign application(s) for patent, inventor's or plant breeder's rights certificate(s), or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent, inventor's or plant breeder's rights certificate(s), or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application No.	Country	Foreign Filing Date (MM/DD/YYYY)	Priority NOT Claimed	Certified Copy Attached	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I hereby appoint Practitioners at Customer Number 36001 as my attorneys and/or agents with full powers (including the powers of appointment, substitution, and revocation) to prosecute this application and any division, continuation, continuation-in-part, reexamination, or reissue thereof, and to transact all business in the U.S. Patent and Trademark Office connected therewith.

**Customer Number for
Practitioner of Record: 32692**


I also appoint the following Practitioners as my attorneys and/or agents with full powers (including the powers of appointment, substitution, and revocation) to prosecute this application and any division, continuation, continuation-in-part, reexamination, or reissue thereof, and to transact all business in the U.S. Patent and Trademark Office connected therewith: None


The mailing address and the telephone number of the above-identified attorneys and/or agents are that of Customer No. 32692.


Inquiries regarding this application can be made to:

Attention: Christopher D. Gram
Office of Intellectual Property Counsel
3M Innovative Properties Company
Telephone No.: (651) 733-1507

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 USC § 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor.
Given Name (first and middle [if any]): Ross M.		Family Name or Surname: Kedl
Inventor's Signature: 		Date: 8/14/03
Residence: Roseville, Minnesota	Country: USA	Citizenship: United States of America
Mailing Address: P.O. Box 33427, St. Paul, Minnesota 55133-3427, USA		

NAME OF JOINT INVENTOR, IF ANY:		<input type="checkbox"/> A petition has been filed for this unsigned inventor.
Given Name (first and middle [if any]): George W.		Family Name or Surname: Griesgraber
Inventor's Signature: 		Date: 8/14/03
Residence: Eagan, Minnesota	Country: USA	Citizenship: United States of America
Mailing Address: P.O. Box 33427, St. Paul, Minnesota 55133-3427, USA		

NAME OF JOINT INVENTOR, IF ANY:		<input type="checkbox"/> A petition has been filed for this unsigned inventor.
Given Name (first and middle [if any]): Isidro Angelo E.		Family Name or Surname: Zarraga
Inventor's Signature: 		Date: Aug. 14, 2003
Residence: Minneapolis, Minnesota	Country: USA	Citizenship: Philippines
Mailing Address: P.O. Box 33427, St. Paul, Minnesota 55133-3427, USA		